



For more information about the booster vaccine, use the QR code below:



For more information about the additional dose for immunocompromised individuals, use the QR code below:



For more information on where to find a COVID-19 vaccine, visit [COVIDVaccineFacts.com](https://www.covidvaccinefacts.com) or click the QR code:



BOOSTERS and ADDITIONAL DOSES

BOOSTER SHOT

Everyone ages 12 years and older should get a COVID-19 booster shot. Over time, the immunity from your initial COVID-19 vaccines can decrease a little. The booster shot helps to “boost” your immune response to restore and enhance protection against COVID-19. Eligible individuals may choose the vaccine type that they originally received or choose a different booster. CDC’s recommendations now allow for this type of mix and match dosing for booster shots; however, the Pfizer or Moderna vaccines are preferred in most [situations](#).

Pfizer and Moderna



COVID-19 vaccine booster shots are recommended for vaccine recipients who completed their initial series at least 5 months ago and who are at least 12 years or older. (Individuals ages 12-17 years should receive a Pfizer booster dose at least 5 months after completing their initial series.)


Johnson & Johnson (Janssen)



ADDITIONAL DOSE for immunocompromised

People who have [moderately to severely compromised immune systems](#) may not mount an adequate immune response to vaccines and therefore may not be protected after their primary series. Individuals age 5 years or older with moderately to severely compromised immune systems at the time of vaccination should get an additional dose of COVID-19 vaccine as part of their primary series, followed by a booster shot (for age 12 years and older only) after the recommended time interval. Use of mRNA vaccines (Pfizer or Moderna) is preferred.

Pfizer and Moderna (4 total doses: 3-dose primary series, then booster)



The same product should be used for all doses of the primary series.

Johnson & Johnson (Janssen) (3 total doses: 2-dose primary series, then booster)



*Any COVID-19 vaccine can be used for the booster dose in people ages 18 years and older, though mRNA vaccines are preferred. For people ages 12-17 years, only Pfizer can be used. People ages 5-11 years should not receive a booster dose. If Moderna used for booster, dose is 50 mcg (0.25 ml)

**Use Pfizer or Moderna. If Moderna used for additional dose, dose is 100 mcg (0.5 ml)